

VYVGART® Hytrulo
(efgartigimod alfa and hyaluronidase-qvfc)
Subcutaneous Injection
180 mg/mL and 2000 U/mL vial

In-Office Subcutaneous Injection Guide

VYVGART Hytrulo—an FDA-approved treatment for adults with CIDP

A first for CIDP—one weekly injection, given under the skin by a healthcare professional, that takes ~30 to 90 seconds.* Get information about what to expect for the in-office injection process: before, during, and after.

*After administering your injection, your healthcare provider will monitor you for allergic reactions for at least 30 minutes.
CIDP=chronic inflammatory demyelinating polyneuropathy; FDA=Food and Drug Administration

What is VYVGART® HYTRULO (efgartigimod alfa and hyaluronidase-qvfc)?

VYVGART HYTRULO is a prescription medicine used for the treatment of adult patients with chronic inflammatory demyelinating polyneuropathy (CIDP)

SELECT IMPORTANT SAFETY INFORMATION

Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see the full Prescribing Information for VYVGART HYTRULO.

VYVGART Hytrulo is one weekly injection



One injection given weekly that takes
~30 to 90 seconds.*

*After administering your injection, your healthcare provider will monitor you for allergic reactions for at least 30 minutes.



A single subcutaneous injection,
delivered under the skin, that **does not**
require vein access.



Given at a **doctor's office, infusion center,**
or at home by a healthcare professional.†

†In some cases, VYVGART Hytrulo may also be given at home by a trained nurse.

SELECT IMPORTANT SAFETY INFORMATION

VYVGART HYTRULO may cause serious side effects, including infection. VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.

Please see Important Safety Information throughout. Please see the full Prescribing Information for VYVGART HYTRULO.

Things to consider before your in-office subcutaneous injection appointment

- Prepare for your visit. You may need to find a ride if you can't drive yourself, and you should plan to get to your appointment at least 15 minutes early
- Consider wearing clothing that's loose around the waist on the day of your injection
- Bring a list of your current medications to your appointment

Keep in mind: how to make the most of the time

VYVGART Hytrulo takes about 30 to 90 seconds to inject, but your healthcare provider will monitor you for allergic reactions for at least 30 minutes after your injection. You may want to bring along something you enjoy doing while you wait.

**Book****Crossword****TV show****Podcast****Game**

SELECT IMPORTANT SAFETY INFORMATION

VYVGART HYTRULO may cause serious side effects, including allergic reactions (hypersensitivity reactions). VYVGART HYTRULO can cause allergic reactions such as rashes, swelling under the skin, and shortness of breath. Hives were also observed in patients treated with VYVGART HYTRULO. Serious allergic reactions, such as trouble breathing and decrease in blood pressure leading to fainting have been reported with efgartigimod alfa-fcab.

Please see Important Safety Information throughout. Please see the full Prescribing Information for [VYVGART HYTRULO](#).

What to expect during your VYVGART Hytrulo subcutaneous injection

- ✓ **Check in**
Once you get to your treatment location, check in with the staff and confirm your insurance information.
- ✓ **Get ready for injection**
An office staff member will help get you ready for your subcutaneous injection. VYVGART Hytrulo may also be prepared for injection at this time.
- ✓ **Receive your injection**
A healthcare professional will inject VYVGART Hytrulo subcutaneously into an area of pinched skin on your abdomen over 30 to 90 seconds. A different area of your abdomen will be used for subsequent injections of VYVGART Hytrulo.
- ✓ **Stay for monitoring**
You will be monitored for allergic reactions for at least 30 minutes after your injection. Consider bringing something to pass the time.
- ✓ **Check out**
Before you leave, make sure to confirm or schedule your next appointment.



Not actual size

Are you enrolled in the VYVGART Patient Support Program?

My VYVGART® Path

There are **2 options** to enroll once your doctor has prescribed you VYVGART Hytrulo:

OPTION 1

Speak with your doctor
and ask to be enrolled in
My VYVGART Path.



OPTION 2

Scan the QR code to
download the patient
enrollment form.



Already enrolled? Reach out to a Nurse Case Manager
for support at **1-833-MY-PATH-1** (1-833-697-2841).

IMPORTANT SAFETY INFORMATION

Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

VYVGART HYTRULO may cause serious side effects, including:

- **Infection.** VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.
- **Allergic Reactions (hypersensitivity reactions).** VYVGART HYTRULO can cause allergic reactions such as rashes, swelling under the skin, and shortness of breath. Hives were also observed in patients treated with VYVGART HYTRULO. Serious allergic reactions, such as trouble breathing and decrease in blood pressure leading to fainting have been reported with efgartigimod alfa-fcab.
- **Infusion-Related Reactions.** VYVGART HYTRULO can cause infusion-related reactions. The most frequent symptoms and signs reported with efgartigimod alfa-fcab were high blood pressure, chills, shivering, and chest, abdominal, and back pain.

Tell your doctor if you have signs or symptoms of an infection, allergic reaction, or infusion-related reaction. These can happen while you are receiving your VYVGART HYTRULO treatment or afterward. Your doctor may need to pause

or stop your treatment. Contact your doctor immediately if you have signs or symptoms of a serious allergic reaction.

Before taking VYVGART HYTRULO, tell your doctor if you:

- take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines,
- have received or are scheduled to receive a vaccine (immunization), or
- have any allergies or medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding.

What are the common side effects of VYVGART HYTRULO?

The most common side effects in efgartigimod-alfa-fcab-treated patients were respiratory tract infection, headache, and urinary tract infection. Additional common side effects with VYVGART HYTRULO are injection site reactions, including rash, redness of the skin, itching sensation, bruising, pain, and hives.

These are not all the possible side effects of VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to the US Food and Drug Administration at 1-800-FDA-1088.

What is VYVGART® HYTRULO (efgartigimod alfa and hyaluronidase-qvfc)?

VYVGART HYTRULO is a prescription medicine used for the treatment of adult patients with chronic inflammatory demyelinating polyneuropathy (CIDP)

Please see the full Prescribing Information for VYVGART HYTRULO and talk to your doctor.